

Results

Prevalencia de disfagia en estudios retrospectivos

Proportions

Random-Effects Model (k = 5)

	Estimate	se	Z	p	CI Lower Bound	CI Upper Bound
Intercept	0.278	0.126	2.20	0.028	0.030	0.526


Note. Tau² Estimator: Restricted Maximum-Likelihood

[3]

Heterogeneity Statistics

Tau	Tau ²	I ²	H ²	R ²	df	Q	p
0.278 (SE= 0.0564)	0.0771	99.62%	261.048	.	4.000	110.565	< .001

Forest Plot



[3]

Publication Bias Assessment

Fail-Safe N Analysis (File Drawer Analysis)

Fail-safe N	p
563.000	< .001

Note. Fail-safe N Calculation Using the Rosenthal Approach


Rank Correlation Test for Funnel Plot Asymmetry

Kendall's Tau	p
1.000	0.017

Regression Test for Funnel Plot Asymmetry

Z	p
7.437	< .001

Funnel Plot



[3]

References

- [1] The jamovi project (2021). *jamovi*. (Version 2.2) [Computer Software]. Retrieved from <https://www.jamovi.org>.
- [2] R Core Team (2021). *R: A Language and environment for statistical computing*. (Version 4.0) [Computer software]. Retrieved from <https://cran.r-project.org>. (R packages retrieved from MRAN snapshot 2021-04-01).
- [3] Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*. [link](#), 36, 1-48.